



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 420 064 A2**

12

EUROPEAN PATENT APPLICATION

21 Application number: 90118198.2

51 Int. Cl.⁵: **C07C 233/23**, C07C 233/06,
C07C 233/74, C07C 233/60,
A61K 31/16

22 Date of filing: 21.09.90

30 Priority: 25.09.89 US 411675

43 Date of publication of application:
03.04.91 Bulletin 91/14

84 Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

71 Applicant: **NORTHWESTERN UNIVERSITY**
633 Clark Street
Evanston Illinois 60208(US)

72 Inventor: **Horn, Alan S.**
Noordwijkweg 3A
NL-9804 RA Noordhorn(NL)
Inventor: **Dubocovich, Margarita L.**
2608 Noyes
Evanston, Illinois 60201(US)

74 Representative: **Patentanwälte Grünecker,**
Kinkeldey, Stockmair & Partner
Maximilianstrasse 58
W-8000 München 22(DE)

54 **Substituted 2-amidotetralins as melatonin agonists and antagonists.**

57 The present invention relates generally to compounds having melatonin receptor activities and in particular to substituted 2-amidotetralin derivatives; to pharmaceutical preparations comprising such compounds; and to methods for using these compounds as therapeutic and diagnostic reagents.


EP 0 420 064 A2

SUBSTITUTED 2-AMIDOTETRALINS AS MELATONIN AGONISTS AND ANTAGONISTS

BACKGROUND OF THE INVENTION

The present invention relates generally to compounds having melatonin receptor activities and in particular to substituted derivatives of 2-amidotetralins, to pharmaceutical preparations comprising these compounds; to methods for using these compounds as therapeutic agents and to methods for using these compounds as diagnostic reagents.

Melatonin (N-acetyl-5-methoxytryptamine) is a hormone synthesized and secreted primarily by the pineal gland, with highest levels occurring during the dark period of a circadian light-dark cycle. The hormone is also found in the retina, and gut. The structure of melatonin is:



Melatonin is involved in the transduction of photoperiodic information and appears to modulate a variety of neural and endocrine functions in vertebrates, including the regulation of reproduction, body weight and metabolism in photoperiodic mammals, the control of circadian rhythms and the modulation of retinal physiology. Retinal melatonin has been implicated in photoreceptor outer segment disc shedding and phagocytosis, in melanosome aggregation in pigment epithelium, and cone photoreceptor retinomotor movement. Of the various physiological processes that have been associated with melatonin, those best substantiated are its effects on sexual maturation, ovarian function, and chronobiological rhythms. See Arendt, J., Oxford Reviews of Reproductive Biology, Vol. 8: 266-320 (1986) and Dubocovich, M.L., FASEB J., 2:2765-2773 (1988).

Exogenous melatonin administration has been found to synchronize circadian rhythms in rats. Cassone, et al., J. Biol. Rhythms, 1:219-229 (1986). In humans, administration of melatonin has been used to treat jet-lag related sleep disturbances, considered to be caused by desynchronization of circadian rhythms. Arendt, et al., Br. Med. J., 292:1170 (1986). Classically, the physiological effects of endogenous melatonin have been antagonized by exposure of animals to light, which inhibits the synthesis of melatonin, or by removal of the pineal gland.

Pharmacological studies have shown that picomolar concentrations of melatonin selectively inhibit the calcium-dependent release of dopamine from rabbit and chicken retina through activation of a site having the pharmacological and functional characteristics of a receptor. Dubocovich, M.L., Nature, 306:782-784 (1983). Dubocovich, M.L., Eur. J. Pharmacol., 105:193-194 (1984). Dubocovich, M.L., J. Pharmacol. Exp. Ther., 234:395-401 (1985). Using the radioiodinated ligand, 2-[¹²⁵I]iodomelatonin, which enhances the ability to detect melatonin receptor sites in vertebrate retina, a pharmacological correlation between the binding site labeled by 2-[¹²⁵I]iodomelatonin and a functional response regulated by melatonin in the chicken and rabbit retina has been demonstrated. Dubocovich, M.L., et al., Proc. Nat'l. Acad. Sci. USA, 84:3916 (1987). Using this radioligand, melatonin binding sites have been localized primarily in the suprachiasmatic nucleus and pars tuberalis/median eminence of mammals, including humans. Reppert, et al., Science, 242:78-81 (1988) and Duncan, et al., Endocrinol., 125:1011-1018 (1989). This is interesting because the melatonin receptor sites localized in the suprachiasmatic nucleus and/or median eminence/pars tuberalis have been suggested to regulate circadian rhythms and reproductive function.

While the radioligand 2-[¹²⁵I]iodomelatonin is a useful probe for the localization and characterization of melatonin receptors, a significant problem in further elucidating the mechanism of action of melatonin is the lack of potent and selective melatonin receptor agonists and antagonists. Such agonists/antagonists could find application not only in the study of melatonin receptor interactions but also in the treatment of conditions possibly affected by melatonin activity, such as depression, jet-lag, disturbances in the sleep-

wakefulness cycle, hypertension, glaucoma, reproduction and neuroendocrine disorders.

Generally, agonists of neurotransmitters and neurohormones are structurally related to the transmitter they mimic, whereas antagonists may be structurally unrelated and quite diverse. The term melatonin agonist is used herein to comprise compounds that mimic melatonin activity, e.g., melatonin inhibition of the calcium-dependent release of [³H]dopamine evoked by electrical stimulation of chicken retina and rabbit retina. To date, all of the known melatonin agonists are derivatives of melatonin itself, e.g., 2-iodomelatonin, 6-chloromelatonin, 6,7-dichloro-2-methyl melatonin, and 8-hydroxymelatonin, and all contain the 5-methoxy indole ring system as an essential moiety. See, Dubocovich, et al., *Proc. Nat'l. Acad. Sci. (USA)*, 84:3916-3918 (1987). Dubocovich, J. *Pharmacol. Exp. Ther.*, 234:395 (1985).


As for melatonin antagonists, previous structure-activity studies on the melatonin receptor led to the suggestion that N-acetyltryptamine analogs lacking the 5-methoxy group might be potential melatonin receptor antagonists. Heward, et al., *Life Sci.*, 17:1167-1178 (1975). Dubocovich, M.L., *Eur. J. Pharmacol.*, 105:193 (1984). Dubocovich, et al., *Proc. Nat'l. Acad. Sci. (USA)*, 184:3916 (1987). Dubocovich, M.L., *J. Pharmacol. Exp. Ther.*, 234:395 (1985). However, while N-acetyltryptamine was reported to antagonize the melatonin-induced lightening of the α -melanocyte stimulating hormone-induced darkening of the frog skin, and to be a competitive melatonin receptor antagonist in chicken retina, it was found to be a partial melatonin agonist in rabbit retina. Heward, C.B., et al., *Life Sci.*, 17:1167 (1975); Dubocovich, M.L., *Eur. J. Pharmacol.*, 105:193 (1984). Dubocovich, M.L., *J. Pharmacol. Exp. Ther.*, 234:395 (1985).

The first compound shown to be a competitive antagonist of mammalian melatonin receptors was luzindole (2-benzyl-N-acetyltryptamine). Dubocovich, M.L., *J. Pharmacol. Exp. Ther.*, 246:902 (1988). Luzindole, in concentrations up to 10 μ M, does not modify either the spontaneous outflow or the calcium-dependent release of [³H]dopamine, but does antagonize the ability of melatonin to inhibit [³H]dopamine release in a competitive fashion. A variety of other compounds have been examined for possible antagonistic activity; however, none have shown the competitive antagonism seen with luzindole.

All of these known melatonin agonists and antagonists retain the 5-melatonin-indole ring system of the natural hormone melatonin. Historically, development of receptor agonists and antagonists which are chemically unrelated to the natural hormone or neurotransmitter has led to selective and potent therapeutic agents. However, there continues to exist a need in the art for such melatonin agonists and antagonists for use in investigating the physiological role of the hormone melatonin and in treating disorders involving disturbances of the melatonin system. The present invention describes a non-indolic class of compounds which surprisingly show high melatonin agonist/antagonist activity.

BRIEF SUMMARY OF THE INVENTION

The compounds of the invention have the general formula:



wherein:


- ⁵⁰ R₁ is hydrogen, halogen, amino, amido, a C₁₋₄ alkyl, alkoxy, or alkoxyaryl;
 - R₂ is hydrogen, hydroxyl, halogen, amino, amido, aryl, mono- or di- C₁₋₄ alkylamino, C₁₋₄ alkylaryl, or alkoxyaryl, or C₁₋₄ alkyl, alkenyl, alkyne, alkoxy;
 - R₃ is hydrogen, aryl, C₁₋₄ alkylaryl, or C₁₋₄ alkyl, alkenyl, or alkyne;
 - R₄ is aryl, C₁₋₄ alkylaryl, or C₁₋₄ alkyl, haloalkyl, or cycloalkyl;
 - ⁵⁵ R₅ is hydrogen, hydroxyl, halogen, oxo, aryl, C₁₋₄ alkylaryl, or C₁₋₄ alkyl;
- wherein aryl substituents of R₂, R₃, R₄, and R₅ may optionally be halogen, hydroxyl, amino, mono- or di- C₁₋₄ alkylamino, or C₁₋₄ alkyl or alkoxy substituted, provided that when R₁ is methoxy, and R₂, R₃, and R₅ are hydrogen, R₄ is not methyl.

In preferred embodiments of the compound, the formulation includes R_5 limited to the 4-position. Most exemplified compounds include H in the R_5 position. Most effective antagonists include phenyl in the R_5 position.

Compounds of the invention contain at least one asymmetric carbon atom, i.e., C_2 and as a result the properties of the compound may, to a greater or lesser extent, be ascribed to either stereoisomer. Thus, the pure enantiomers, as well as the racemic mixtures thereof, are within the scope of the invention. For compounds containing two asymmetric carbon atoms, and therefore occurring as cis or trans isomers, it is understood that the invention encompasses both isomers.

Preferred compounds include: 8-methoxy-2-propionamido-tetralin; 2-chloroacetamido-tetralin; 8-methoxy-2-n-butyramido-tetralin; 8-methoxy-2-cyclopropanecarbonylamido-tetralin; 8-methoxy-2-chloroacetamido-tetralin; 4-phenyl-2-acetamido-tetralin; 4-phenyl-2-propionamido-tetralin; 4-benzyl-2-acetamido-tetralin; 4-phenyl-2-chloroacetamido-tetralin; and 4-benzyl-2-propionamido-tetralin.

Compounds of the invention exhibit strong melatonin receptor activity and are expected to be useful in methods for treating disorders which arise from a disturbed functioning of the melatonin system. Thus, also provided by the invention are methods for treatment of disorders associated with abnormal melatonin activity in an organism wherein a therapeutically effective amount of a composition is administered comprising a compound of the formula:



wherein:

- R_1 is hydrogen, halogen, amino, amido, a C_{1-4} alkyl, alkoxy, or alkoxyaryl;
 - R_2 is hydrogen, hydroxyl, halogen, amino, amido, aryl, mono- or di- C_{1-4} alkylamino, C_{1-4} alkylaryl, or alkoxyaryl, or C_{1-4} alkyl, alkenyl, alkynyl, alkoxy;
 - R_3 is hydrogen, aryl, C_{1-4} alkylaryl, or C_{1-4} alkyl, alkenyl, or alkynyl;
 - R_4 is aryl, C_{1-4} alkylaryl, or C_{1-4} alkyl, haloalkyl, or cycloalkyl;
 - R_5 is hydrogen, hydroxyl, halogen, oxo, aryl, C_{1-4} alkylaryl, or C_{1-4} alkyl;
- wherein aryl substituents of R_2 , R_3 , R_4 , and R_5 may optionally be halogen, hydroxyl, amino, mono- or di- C_{1-4} alkylamino, or C_{1-4} alkyl or alkoxy substituted.

Preferred compounds for this method include those where R_1 is hydrogen or alkoxy, where R_2 is hydrogen or alkoxy, where said alkoxy is preferably methyl, where R_3 is preferably hydrogen, and where R_4 is preferably methyl, chloromethyl, ethyl, propyl, cyclopropyl, n-butyl, iso-butyl, phenyl, or benzyl; and/or where R_5 is phenyl or benzyl. Specifically, preferred compounds are those having melatonin receptor activity including: 8-methoxy-2-acetamido-tetralin; 8-methoxy-2-propionamido-tetralin; 2-chloroacetamido-tetralin; 8-methoxy-2-n-butyramido-tetralin; 8-methoxy-2-cyclopropanecarbonylamido-tetralin; 8-methoxy-2-chloroacetamido-tetralin; 4-phenyl-2-acetamido-tetralin; 4-phenyl-2-propionamido-tetralin; 4-benzyl-2-acetamido-tetralin; 4-phenyl-2-chloroacetamido-tetralin; and 4-benzyl-2-propionamido-tetralin.

Presently preferred compounds for use in a method for blocking melatonin function by administering a melatonin antagonist include: 4-phenyl-2-acetamido-tetralin; 4-phenyl-2-propionamido-tetralin; and 4-phenyl-2-chloroacetamido-tetralin. Presently preferred compounds for use in a method for mimicking melatonin function by administering a melatonin agonist include: 8-methoxy-2-acetamido-tetralin; 8-methoxy-2-propionamido-tetralin; 2-chloroacetamido-tetralin; 8-methoxy-2-n-butyramido-tetralin; 8-methoxy-2-cyclopropanecarbonylamido-tetralin; 8-methoxy-2-chloroacetamido-tetralin; 4-benzyl-2-acetamido-tetralin; and 4-benzyl-2-propionamido-tetralin.

Also provided by the invention are pharmaceutical compositions comprising pharmaceutically effective amounts of one or more compounds of the invention in combination with an inert pharmaceutical carrier, as well as methods for producing a melatonin-like action, or alternatively blocking the effects of melatonin, by administering these pharmaceutical compositions. Generally, where it is desirable to inhibit melatonin function, those compounds of the invention having melatonin antagonist activity are administered; where it is desirable to mimic melatonin function, those compounds having melatonin agonist activity are administered.

Some compounds (i.e., partial agonists) may show either agonist or antagonist activity depending not only on the activity of the melatonergic system at the time of administration, but also upon both the dose and route of administration. The route of administration can be any that produces the desired effect and includes for example oral, intra-venous, intra-muscular, and topical applications, including, e.g., time release transdermal patches.

In one preferred embodiment of the present invention, a therapeutically effective amount of one or more of the compounds of the invention can be administered to treat chronobiological disorders, such as, seasonal affective disorder (SAD), sleep disorders, and symptoms, such as drowsiness and fatigue, associated with disturbances in sleep/wake cycles (e.g., jet-lag, workers on night shifts, etc.), or to treat glaucoma by lowering intra-ocular pressure. Therapeutically effective amounts of one or more compounds of the invention can be administered to treat various psychiatric disorders related to altered melatonin function or influenced by melatonin and circadian rhythms, e.g., affective disorders (mania and depression), alcoholism, stress, aging (dementia such as Alzheimer's disease), epilepsy, convulsions, anxiety, tremors induced by idiopathic parkinsonism, and adventitious movements induced by L-dopa movements; or to treat various neuroendocrine disorders related to altered melatonin function or influenced by melatonin and biological rhythms (e.g., peptic ulceration, psoriasis, hair tonic, and body weight) particularly relating to regulation of reproductive maturation and function e.g., idiopathic delayed puberty, sudden infant death (SID), premature labor, infertility, antifertility, premenstrual syndrome, and sexual dysfunction (i.e., impotence and decreased orgasmic activity). In addition, compounds of the invention can be used alone, or in combination, in animals to manipulate breeding cycles, body weight, coat color and oviposition and can be used as ovulation or sperm production inhibitors to control the population of susceptible hosts, including insects, birds, and animals.

Also provided by the invention is a method for localizing melatonin receptors in for example, a tissue sample, wherein the sample is incubated with a compound of the invention and wherein the compound is labelled with a marker, the marker is detected and the melatonin receptors in the sample are thereby localized.

Other aspects and advantages of the present invention will be apparent upon consideration of the following detailed description thereof which includes numerous illustrative examples of the practice of the invention.

DETAILED DESCRIPTION OF THE INVENTION

The following examples 1 through 26 are intended to be illustrative of practice of the present invention rather than limiting thereon. Examples 1-22 describe the synthesis of the compounds of the invention. Examples 23-25 provide the results obtained from the pharmacological testing of the compounds of the invention. Specifically, example 23 provides the results obtained with the compounds of the invention in an assay that measures the inhibition of 2-[¹²⁵I]iodomelatonin binding using chicken retinal membranes; example 24 provides the results obtained in an assay to ascertain which test compounds mimic the melatonin-induced inhibition of the release of [³H]dopamine from rabbit retina membranes; example 25 provides the results obtained in an assay to measure the potential antagonistic activity of the test compounds during the melatonin-induced inhibition of [³H]dopamine release from rabbit retina.

Example 26 relates to pharmaceutical preparations comprising the compounds of the invention, to methods for using these compounds as therapeutic agents in treating disorders associated with abnormal melatonin activity, and to methods for using the compounds as diagnostic reagents.

SYNTHESIS OF COMPOUNDS OF THE INVENTION

The compounds of the invention are synthesized as described in Examples 1-22 below. Table 1 provides: the example number which describes the synthesis of the compound; the reference number and name of each compound; and the substituents: R₁, R₂, R₃, R₄, and R₅ for each compound. The general formula for the compounds synthesized is given below.




TABLE 1

EXAMPLE	COMPOUND AND REFERENCE NUMBER	SUBSTITUENTS			
		R ₁	R ₂ N(R ₃)-C(O)-R ₄	R ₅	
2	AH001 8-methoxy-2-acetamido-tetralin	OCH ₃	H NHC(O)CH ₃	H	
1	AH002 8-methoxy-2-propionamido-tetralin	OCH ₃	H NHC(O)C ₂ H ₅	H	
10	AH005 2-acetamido-tetralin	H	H NHC(O)CH ₃	H	
11	AH006 2-propionamido-tetralin	H	H NHC(O)C ₂ H ₅	H	
3	AH007 8-methoxy-2-benzoylamido-tetralin	OCH ₃	H NHC(O)benzyl	H	
8	AH008 7-methoxy-2-acetamido-tetralin	H	OCH ₃ NHC(O)CH ₃	H	
12	AH010 2-phenylacetamido-tetralin	H	H NHC(O)benzyl	H	
13	AH011 2-chloroacetamido-tetralin	H	H NHC(O)CH ₂ Cl	H	
4	AH013 8-methoxy-2-n-butylamido-tetralin	OCH ₃	H NHC(O)C ₃ H ₇	H	
5	AH014 8-methoxy-2-iso-butylamido-tetralin	OCH ₃	H NHC(O)CH(CH ₃) ₂	H	
6	AH015 8-methoxy-2-cyclopropanecarbonylamido-tetralin	OCH ₃	H NHC(O)C ₃ H ₅	H	
7	AH016 8-methoxy-2-phenylacetamido-tetralin	OCH ₃	H NHC(O)benzyl	H	

TABLE 1 - Continued

EXAMPLE	COMPOUND AND REFERENCE NUMBER	R ₁	SUBSTITUENTS		R ₅
			R ₂	$\overline{\text{N(R}_3\text{)-C(O)-R}_4}$	
16	AH017 8-methoxy-2-chloroacetamido-tetralin	OCH ₃	H	NHC(O)CH ₂ Cl	H
9	AH018 5-methoxy-2-acetamido-tetralin	H	OCH ₃	NHC(O)CH ₃	H
14	AH019 2-benzoylamido-tetralin	H	H	NHC(O)phenyl	H
15	AH020 2-butyramido-tetralin	H	H	NHC(O)C ₃ H ₇	H
17	AH023 4-phenyl-2-acetamido-tetralin	H	H	NHC(O)CH ₃	phenyl
18	AH024 4-phenyl-2-propionamido-tetralin	H	H	NHC(O)C ₂ H ₅	phenyl
20	AH025 4-benzyl-2-acetamido-tetralin	H	H	NHC(O)CH ₃	benzyl
19	AH026 4-phenyl-2-chloroacetamido-tetralin	H	H	NHC(O)CH ₂ Cl	phenyl
21	AH027 4-benzyl-2-propionamido-tetralin	H	H	NHC(O)C ₂ H ₅	benzyl
20d	AH028 4-benzyl-2-acetamido-1-tetralone	H	H	NHC(O)CH ₃	benzyl
22	AH029 4-benzyl-2-chloroacetamido-tetralin	H	H	NHC(O)CH ₂ Cl	benzyl

The three methods used for synthesis of the compounds of the invention are as follows.

5 Method 1:

The appropriately substituted 2-tetralone is converted to the corresponding 2-aminotetralin by condensation with benzylamine, catalytic reduction over platinum, and subsequent debenzylation over palladium-charcoal. The resulting primary amine is then reacted with the appropriate acyl chloride or acid
 10 anhydride to yield the desired amide. Horn, A.S., et al., J. Med. Chem., 21:825 (1978). The following amides were synthesized utilizing the indicated reactants:

Amide	Substituted 2-tetralone	Acid anhydride/Acyl Chloride
AH 001	8-methoxy-2-tetralone	acetic anhydride
AH 002	8-methoxy-2-tetralone	propionic anhydride
AH 005	2-tetralone	acetic anhydride
AH 006	2-tetralone	propionic anhydride
AH 007	8-methoxy-2-tetralone	benzoyl chloride
AH 008	7-methoxy-2-tetralone	acetic anhydride
AH 010	2-tetralone	phenylacetyl chloride
AH 011	2-tetralone	chloroacetyl chloride
AH 013	8-methoxy-2-tetralone	butyric anhydride
AH 014	8-methoxy-2-tetralone	isobutyryl chloride
AH 015	8-methoxy-2-tetralone	cyclopropanecarbonyl chloride
AH 016	8-methoxy-2-tetralone	phenylacetyl chloride
AH 017	8-methoxy-2-tetralone	chloroacetyl chloride
AH 018	5-methoxy-2-tetralone	acetic anhydride
AH 019	2-tetralone	benzoyl chloride
AH 020	2-tetralone	butyric anhydride
AH 023	4-phenyl-2-tetralone	acetic anhydride
AH 024	4-phenyl-2-tetralone	propionic anhydride
AH 026	4-phenyl-2-tetralone	chloroacetyl chloride

METHOD 1

5


10

15

20

25

30

**Method 2:**






The appropriately substituted 1-tetralone is converted to its oxime and this compound is treated with p-toluenesulfonyl chloride to yield the O-(p-tolylsulfonyl)oxime. Treatment of the latter compound with sodium ethoxide in benzene produces the keto-amine via the Neber rearrangement. Acylation of this derivative with the appropriate acyl halide or acid anhydride yields the corresponding keto-amide. Catalytic reduction of this intermediate over palladium-charcoal yields the desired end product. Horn, A.S., et al., Eur. J. Med. Chem., 23:325 (1988). Weinstock, J., et al., J. Med. Chem., 29:1615 (1986). The following amides were synthesized according to method 2 utilizing the reactants specified.

45

50

Amide	Substituted 1-tetralone	Acid anhydride/Acyl chloride
AH 025	4-benzyl-1-tetralone	acetic anhydride
AH 027	4-benzyl-1-tetralone	propionic anhydride
AH 028	4-benzyl-1-tetralone	acetic anhydride
AH 029	4-benzyl-1-tetralone	chloroacetyl chloride

55

METHOD 2

50 The synthesis of analogues having various R_3 substituents is achieved through standard variations of the methods described in the examples below, i.e., using one of the two general methods 1 or 2 to obtain the appropriately substituted intermediate which is then acylated using the appropriate R_4 substituted acid chloride or anhydride. The synthesis of analogues having various R_4 substituents is shown in general method 3.

55

Method 3:

The appropriately substituted 2-tetralone is treated with the corresponding R_3 substituted amine; the

resulting intermediate is reduced; and the final product is obtained by acylation with the appropriate acid chloride or anhydride. Hacksell, U., et al., J. Med. Chem., 22:1469 (1979).

METHOD 3

5

10



15

20

25

30

35



EXAMPLE 1

40

Preparation of 8-methoxy-2-propionamidotetralin [AH 002]:

To a solution of 8-methoxy-2-tetralone (5.30 g, 30.1 mmol) in 50 ml of benzene, benzylamine (4.30 g, 40 mmol) and p-toluenesulfonic acid (0.1 g) were added. The mixture was then refluxed for 1 hr under an atmosphere of nitrogen gas using a Dean and Stark apparatus. The benzene and other volatiles were removed under reduced pressure and 45 ml of absolute alcohol was added to the residue. After transfer to a Parr hydrogenation flask, platinum oxide (50 mg) was added and the mixture was hydrogenated for 2.5 hr under a hydrogen pressure of 2 atmospheres. After filtration of the mixture and removal of the ethanol under reduced pressure, a brown oil was obtained. This oil was diluted with ethyl acetate (25 ml) and then a solution of ether/HCl was added dropwise to yield the HCl salt of 8-methoxy-2-benzylaminotetralin (7.20 g, 79%). Three grams (9.9 mmol) of this salt was converted to the free base by treatment with a solution of sodium carbonate. The base was extracted into ether which was then dried and evaporated under reduced pressure to yield a brown oil. The latter was dissolved in absolute alcohol (60 ml) and hydrogenated in a Parr flask over palladium-charcoal (1.6 g) for 1 hr at 2-atmospheres. Following an isolation procedure similar to that described above, 8-methoxy-2-aminotetralin HCl (1.17 g, 55%) was isolated.

m.p. 281-282 °C.
 I.R. (KBr) 3160 cm^{-1} 2420 cm^{-1} (NH_3^+) 2060 cm^{-1} (NH_3^+).

M.S. (chemical ionization, NH_3 gas) m/e 178 ($M + 1$).

- To a solution of 8-methoxy-2-aminotetralin HCl (0.30 g, 1.40 mmol) in ethyl acetate (15 ml) water (5 ml) and sodium acetate (0.65 g), propionic anhydride (1.15 ml) was added and the mixture was stirred well for 6 hr. Water (10 ml) was then added and after shaking the aqueous and organic layers were separated. The aqueous layer was extracted with ethylacetate and the combined ethyl acetate layers were washed with saturated solutions of sodium bicarbonate and sodium chloride and then dried (MgSO_4) and evaporated under reduced pressure to yield a white solid. Recrystallization (methanol/ethyl acetate) yielded 0.10 g (31%) of a white crystalline material.
- m.p. 151-152 °C.
- 10 I.R. (KBr) 3315 cm^{-1} (NH) 1640 cm^{-1} (C=O).
- M.S. (chemical ionization, NH_3 gas) m/e 234 ($M + 1$).

EXAMPLE 2

15

Preparation of 8-methoxy 2-acetamido-tetralin [AH 001]:

- 20 Using a procedure similar to that described in Example 1, 0.30 g of 8-methoxy-2-amino-tetralin yielded 0.23 g (69%) of 8-methoxy-2-acetamido-tetralin following treatment with acetic anhydride.
- m.p. 156-157 °C.
- I.R. (KBr) 1640 cm^{-1} (C=O).
- M.S. (chemical ionization, NH_3 gas) m/e 220 ($M + 1$).

25

EXAMPLE 3

30

Preparation of 8-methoxy-2-benzoylamido-tetralin [AH 007]:

- Using a procedure similar to that described in Example 1, 0.40 g of 8-methoxy-2-amino-tetralin yielded 0.44 g (84%) of 8-methoxy-2-benzoylamido-tetralin following treatment with benzoyl chloride.
- 35 m.p. 173-175 °C.
- I.R. (KBr) 1625 cm^{-1} (C=O).
- M.S. (chemical ionization, NH_3 gas) m/e 282 ($M + 1$).

40

EXAMPLE 4

Preparation of 8-methoxy-2-n-butyramidotetralin [AH 013]:

45

- Using a procedure similar to that described in Example 1, 0.22 g of 8-methoxy-2-aminotetralin yielded 0.16 g (63%) of 8-methoxy-2-n-butyramidotetralin following treatment with n-butyric anhydride.
- m.p. 139-140 °C.
- I.R. (KBr) 3300 cm^{-1} (NH) 1630 cm^{-1} (C=O).
- 50 M.S. (chemical ionization, NH_3 gas) m/e 248 ($M + 1$).

EXAMPLE 5

55

Preparation of 8-methoxy-2-iso-butyramido-tetralin [AH 014]:

Using a procedure similar to that described in Example 1, 0.22 g of 8-methoxy-2-amino-tetralin yielded 0.18 g (71%) of 8-methoxy-2-iso-butyramido-tetralin following treatment with iso-butyryl chloride.

m.p. 167-168° C.

I.R. (KBr) 1635 cm^{-1} (C=O).

5 M.S. (chemical ionization, NH_3 gas) m/e 248 (M + 1).

EXAMPLE 6

10

Preparation of 8-methoxy-2-cyclopropanecarbonylamido-tetralin [AH 015]:

Using a procedure similar to that described in Example 1, 0.20 g of 8-methoxy-2-amino-tetralin yielded 0.19 g (83%) of 8-methoxy-2-cyclopropane-carbonylamido-tetralin following treatment with cyclopropanecarbonyl chloride.

m.p. 185-187° C.

I.R. (KBr) 1630 cm^{-1} (C=O).

M.S. (chemical ionization, NH_3 gas) m/e 246 (M + 1).

20

EXAMPLE 7

25

Preparation of 8-methoxy-2-phenylacetamido-tetralin [AH016]:

Using a procedure similar to that described in Example 1, 0.24 g 8-methoxy-2-amino-tetralin yielded 0.20 g (61%) of 8-methoxy-2-phenylacetamido-tetralin following treatment with phenylacetyl chloride.

30 m.p. 154-156° C.

I.R. (KBr) 1630 cm^{-1} (C=O).

M.S. (chemical ionization, NH_3 gas) m/e 296 (M + 1).

35

EXAMPLE 8

Preparation of 7-methoxy-2-acetamido-tetralin [AH 008]:

40

Using a procedure similar to that described in Example 1, 0.40 g of 7-methoxy-2-amino-tetralin yielded 0.21 g (51%) of 7-methoxy-2-acetamido-tetralin following treatment with acetic anhydride.

m.p. 105-108° C.

I.R. (KBr) 1625 cm^{-1} (C=O).

45 M.S. (chemical ionization, NH_3 gas) m/e 220 (M + 1).

EXAMPLE 9

50

Preparation of 5-methoxy-2-acetamido-tetralin [AH 018]:

Using a procedure similar to that described in Example 1, 0.30 g of 5-methoxy-2-amino-tetralin yielded 0.06 g (20%) of 5-methoxy-2-acetamido tetralin following treatment with acetic anhydride.

m.p. not determined

I.R. (KBr) 1625 cm^{-1} (C=O).

M.S. (chemical ionization, NH_3 gas) m/e 220 (M + 1).

EXAMPLE 10**5 Preparation of 2-acetamido-tetralin [AH 005]:**

Using a procedure similar to that described in Example 1, 1.50 g of 2-amino-tetralin yielded 1.01 g (65%) of 2-acetamido-tetralin following treatment with acetic anhydride.

m.p. 109-111 °C.

10 I.R. (KBr) 1630 cm⁻¹ (C=O).

M.S. (chemical ionization, NH₃ gas) m/e 190 (M + 1).

EXAMPLE 11

15

Preparation of 2-propionamido-tetralin [AH 006]:

20 Using a procedure similar to that described in Example 1, 0.26 g of 2-amino-tetralin yielded 0.25 g (86%) of 2-propionamido-tetralin following treatment with propionic anhydride.

m.p. 99-101 °C.

I.R. (KBr) 1635 cm⁻¹ (C=O).

M.S. (chemical ionization, NH₃ gas) m/e 204 (M + 1).

25

EXAMPLE 12

30

Preparation of 2-phenylacetamido-tetralin [AH 010]:

Using a procedure similar to that described in Example 1, 0.45 g of 2-amino-tetralin yielded 0.47 g (72%) of 2-phenylacetamido-tetralin following treatment with phenylacetyl chloride.

35 m.p. 119-121 °C.

I.R. (KBr) 1625 cm⁻¹ (C=O).

M.S. (chemical ionization, NH₃ gas) m/e 266 (M + 1).

40

EXAMPLE 13**Preparation of 2-chloroacetamido-tetralin [AH 011]:**

45

Using a procedure similar to that described in Example 1, 0.38 g of 2-amino-tetralin yielded 0.26 g (56%) of 2-chloroacetamido-tetralin following treatment with chloroacetyl chloride.

m.p. 131-133 °C.

I.R. (KBr) 1650 cm⁻¹ (C=O).

50 M.S. (chemical ionization, NH₃ gas) m/e 224 (M + 1).

EXAMPLE 14

55

Preparation of 2-benzoylamido-tetralin [AH 019]:

Using a procedure similar to that described in Example 1, 0.40 g of 2-amino-tetralin yielded 0.38 g (69%) of 2-benzoylamido-tetralin following treatment with benzoyl chloride.

m.p. 154-156° C.

I.R. (KBr) 1625 cm⁻¹ (C=O).

5 M.S. (chemical ionization, NH₃ gas) m/e 252 (M+1).

EXAMPLE 15

10

Preparation of 2-butyramido-tetralin [AH 020]:

Using a procedure similar to that described in Example 1, 0.24 g of 2-amino-tetralin yielded 0.23 g (81%) of 2-butyramido-tetralin following treatment with butyric anhydride.

15 m.p. 78-81° C.

I.R. (KBr) 1630 cm⁻¹ (C=O).

M.S. (chemical ionization, NH₃ gas) m/e 218 (M+1).

20

EXAMPLE 16

Preparation of 8-methoxy-2-chloroacetamidotetralin [AH 017]:

To a solution of 8-methoxy-2-amino-tetralin HCl (0.16 g, 0.75 mmol) in dichloromethane (10 ml) and in sodium hydroxyde (0.2 g) in water (2 ml), was added chloroacetyl chloride (2 ml) and the mixture was stirred for 6 hr. The reaction was then poured into water (25 ml) and the organic layer was separated. The aqueous layer was extracted with dichloromethane and the combined organic layers were then shaken with a solution of sodium bicarbonate, dried over MgSO₄ and then evaporated to yield a white solid. After recrystallization (acetone/hexane) 0.12 g (63%) of 8-methoxy-2-chloroacetamidotetralin was obtained.

30 m.p. 136-139° C.

I.R. (KBr) 3270 cm⁻¹ (NH) 1640 cm⁻¹ (C=O).

35 M.S. (chemical ionization, NH₃ gas) m/e 254 (M+1).

EXAMPLE 17

40

Preparation of 4-phenyl-2-acetamidotetralin [AH 023]:

a. 4-Phenyl-2-benzylaminotetraline HCl

To a solution of 4.20 g (19.1 mmol) of 4-phenyl-2-tetralone (Fine and Stern, J. Org. Chem., 32:4132 (1967)) in 35 ml of benzene, were added, 1.95 g (18.3 mmol) of benzylamine and a crystal of p-toluenesulfonic acid. The reaction mixture was refluxed under an atmosphere of nitrogen for one hour with continual removal of water via a Dean and Stark trap. Removal of all volatiles under reduced pressure yielded an oil (5.75 g). This oil was dissolved in 50 ml of a mixture of tetrahydrofuran and methanol (15:1). The pH was then adjusted to between 4 and 5 with dry ether-HCl. The precipitate was removed by filtration and 1550 mg of sodiumcyanoborohydride was added to the resulting solution. This mixture was stirred and warmed to 40° C and kept at this temperature for 20 hrs. Occasional adjustments of the pH were necessary. The course of the reaction was followed by the disappearance of the imine peak at 1625 cm⁻¹ in the I.R. spectrum.

55 The reaction mixture was poured into water and made basic with sodium carbonate. After extraction with ether, drying over magnesium sulfate and decolorization with charcoal, the organic extracts were

evaporated under reduced pressure to yield an oil (5.0 g). Conversion to an HCl salt yielded 2.91 g (44%) of 4-phenyl-2-benzylaminotetralin HCl. Recrystallization gave a white crystalline product.

m.p. 204.5-206° C.

I.R. (KBr) 2920-2300 cm^{-1} (NH_3^+).

5 M.S. (chemical ionization, NH_3 gas) m/e 314 ($M + 1$).

b. 4-Phenyl-2-aminotetralin HCl:

10 Three hundred eighty mg (0.82 mmol) of the free base of 4-phenyl-2-benzylaminotetralin was dissolved in 10 ml of ethanol and to which was added 250 mg Pd/C (10%). This mixture was shaken in a Parr apparatus under a pressure of 3 atm. of hydrogen at 35° C for 3 hrs and for a further 18 hrs at room temperature. After removal of the catalyst and evaporation of the organic solvent, an oil was obtained. Conversion to an HCl salt gave 140 mg (40%) of 4-phenyl-2-aminotetralin HCl.

15 m.p. 237-237.7° C.

I.R. (KBr) 3100-2400 cm^{-1} (NH_3^+).

M.S. (chemical ionization, NH_3 gas) m/e 224 ($M + 1$).

20 c. 4-Phenyl-2-acetamidotetralin

To a solution of 180 mg (0.80 mmol) of 4-phenyl-2-aminotetralin HCl in 3.5 ml ethyl acetate and 2.0 ml water, was added 360 mg of sodium acetate. To this mixture 0.6 ml of acetic anhydride was then added and the mixture was stirred for 24 hrs. The addition of 2 ml of water resulted in the formation of two layers.

25 After separation, the water layer was extracted twice with ethyl acetate. The combined organic layers were washed 3 times with a saturated solution of sodium bicarbonate and once with a saturated solution of sodium chloride. After drying over magnesium sulfate, the organic solution was evaporated under reduced pressure to yield 4-phenyl-2-acetamidotetralin as a white solid (170 mg, 79%).

m.p. 176-177.6° C.

30 I.R. (KBr) 3270 cm^{-1} (NH) 1630 and 1550 cm^{-1} (C=O).

M.S. (chemical ionization, NH_3 gas) m/e 266 ($M + 1$).

EXAMPLE 18

35

Preparation of 4-phenyl-2-propionamido-tetralin [AH 024]:

40 Using a procedure similar to that described in Example 17, 100 mg of 4-phenyl-2-amino-tetralin yielded 40 mg (32%) of 4-phenyl-2-propionamido tetralin following treatment with propionic anhydride.

m.p. 144-146° C.

I.R. (KBr) 1640 cm^{-1} (C=O).

M.S. (chemical ionization, NH_3 gas) m/e 280 ($M + 1$).

45

EXAMPLE 19

50

Preparation of 4-phenyl-2-chloroacetamido-tetralin [AH 026]:

Using a procedure similar to that described in Example 17, 130 mg of 4-phenyl-2-amino-tetralin yielded 10 mg (6%) of 4-phenyl-2-chloroacetamido-tetralin following treatment with chloroacetyl chloride.

55 m.p. 189-190° C.

I.R. (KBr) 1640 cm^{-1} (C=O).

M.S. (chemical ionization, NH_3 gas) m/e 300 ($M + 1$).

EXAMPLE 205 Preparation of 4-benzyl-2-acetamidotetralin [AH 025]:a. 4-Benzyl-1-tetralon oxime

10 To 63.3 g (0.7 mol) of 4-benzyl-1-tetralone (Beyer, Chem. Berichte, 70:1101 (1937)) dissolved in 500 ml ethanol, was added 80 g (1.15 mol) of hydroxylamine HCl and 80 g (0.98 mol) of sodium acetate. The mixture was refluxed for 2.5 hrs and then poured into 1.2 L water. After filtration, the white solid was washed with ice-cold water until the pH of the filtrate was 6-7. The product was recrystallized from ethanol yielding 57.5 g (85%).

15 m.p. 142.6-143.5 °C.

I.R. (KBr) 3200 cm⁻¹ (OH), 1590 cm⁻¹ (C=N).

M.S. (chemical ionization, NH₃ gas) m/e 252 (M+1).

20 b. 4-Benzyl-1-tetralon-O-(p-toluenesulfonyl)oxime

The above oxime (48.5 g, 0.19 mol) was dissolved in 190 ml dry pyridine and then cooled to 0 °C. A solution of 150 g (0.78 mol) of p-toluenesulfonyl chloride in 230 ml dry pyridine was added dropwise over 45 min and then left to stir for a further 18 hrs at 0 °C. The orange colored reaction mixture was poured into 25 1.5 L ice-water and the resulting yellowish precipitate was filtered and washed with water and ice-cold ethanol. The product was recrystallized from ethyl acetate to yield 65 g (84%).

m.p. 159-160 °C.

I.R. (KBr) 1590 cm⁻¹ (Ar, C=N), 1370 cm⁻¹, 1180 cm⁻¹ (SO₂).

M.S. (chemical ionization, NH₃ gas) m/e 406 (M+1).

30

c. 2-Amino-4-benzyl-1-tetralone HCl

To a solution of potassium t-butoxide (6.08 g, 54.2 mmol) dissolved in 60 ml of "super-dry" ethanol at 35 0 °C under nitrogen, was added, dropwise over 45 min, a solution of 20 g (49.4 mmol) of the above tosyloxime in 360 ml dry toluene. The reaction mixture was stirred for 2 hrs at 0-4 °C and for 2 hrs at 10-15 °C. After filtration, the filtrate was acidified with 14 ml 36% hydrochloric acid and stirred for 16 hrs at room temperature. The product was filtered and washed twice with 25 ml cold acetone. The filtrate was reduced in volume to obtain a further fraction of the product.

40 Total yield 12.65 g (89%).

m.p. the substance decomposed above 180 °C.

I.R. (KBr) 3180 cm⁻¹ (NH₃⁺) 1680 cm⁻¹ (C=O).

M.S. (chemical ionization, NH₃ gas) m/e 252 (M+1).

AH 028

45

d. 2-Acetamido-4-benzyl-1-tetralone

50 Eight g (27.8 mmol) of the above keto-amine HCl was dissolved in ethyl acetate. To this solution was added 20.7 g (0.15 mol) sodium acetate, 25 ml water and 16.7 ml acetic anhydride. This mixture was stirred for 2 hrs at room temperature. Fifty ml of water were added and after 30 min of stirring, the organic layer was separated and washed with 50 ml of water and 100 ml of saturated NaCl solution. After evaporation under reduced pressure, the resulting oil was dissolved in 40 ml of toluene and refluxed for 30 min using a Dean and Stark trap. Removal of the solvent gave an oil which slowly crystallized giving 6.5 g (80%) of a 55 white solid. Recrystallization from ethanol yielded the product as a white crystalline material.

m.p. 142-143 °C.

I.R. (KBr) 3380 cm⁻¹ (NH) 1695 cm⁻¹ (C=O) 1630 cm⁻¹ (C=O, amide).

M.S. (chemical ionization, NH₃ gas) m/e 294 (M+1).

e. 4-Benzyl-2-acetamidotetralin

Two g (6.8 mmol) of the above keto-amide was dissolved in 30 ml of acetic acid. To this solution, were added 1.4 ml 70% HClO₄ and 340 mg 10% Pd/C. Hydrogenation was carried out for 16 hrs at room temperature under 3 atmospheres of hydrogen. After filtration, the filtrate was evaporated to yield an oil. This oil was dissolved in 150 ml of water and then neutralized with sodium bicarbonate. This solution was extracted twice with dichloromethane (150 ml). The dichloromethane extracts were then shaken with a saturated solution of sodium chloride. After drying over magnesium sulfate, the solvent was removed to yield an oil. This oil was dissolved in dry ether. This resulting solution yielded a white crystalline product, 1.33 g (70%).
 m.p. 168-168.5° C.
 I.R. (KBr) 3280 cm⁻¹ (NH) 1625 cm⁻¹ (C=O).
 M.S. (chemical ionization, NH₃ gas) m/e 280 (M + 1).

EXAMPLE 21Preparation of 4-benzyl-2-propionamido-tetralin [AH 027]:

Using a procedure similar to that described in Example 20d, 500 mg of 2-propionamido-4-benzyl-1-tetralone yielded 180 mg (38%) of 4-benzyl-2-propionamido-tetralin following catalytic reduction.
 m.p. 148-149° C.
 I.R. (KBr) 1650 cm⁻¹ (C=O).
 M.S. (chemical ionization, NH₃ gas) m/e 294 (M + 1).

EXAMPLE 22Preparation of 4-benzyl-2-chloroacetamido-tetralin [AH 029]:

Using a procedure similar to that described in Example 20d, 500 mg of 2-chloroacetamido-4-benzyl-1-tetralone yielded 130 mg (28%) of 4-benzyl-2-chloroacetamido-tetralin following catalytic reduction.
 m.p. 151-152° C.
 I.R. (KBr) 1665 cm⁻¹ (C=O).
 M.S. (chemical ionization, NH₃ gas) m/e 314 (M + 1).

PHARMACOLOGICAL STUDIES

Two types of in vitro tests, specific for melatonin-like activity, were used for evaluating the pharmacological activities of the synthesized compounds and are described below in Examples 23-25.

The first test, described in Example 23, measures the ability of a test compound to inhibit 2-[¹²⁵I]-iodomelatonin binding to a site exhibiting the pharmacological characteristics of a melatonin receptor in chicken retinal membranes. If a compound shows high activity (i.e., a low IC₅₀) in inhibiting the binding of 2-[¹²⁵I]iodomelatonin to the melatonin binding site, this demonstrates the compound's ability to interact with the melatonin binding site; however, such activity does not establish whether the test compound is an agonist or an antagonist. Examples 24 and 25, respectively, describe tests to establish agonist or antagonist activity.

The Example 24 assay ascertains whether a test compound mimics the melatonin-induced inhibition of the release of [³H]dopamine from rabbit retina membranes. In this assay, the test compound is added before the second period of stimulation (S₂) and its effect is compared to that of melatonin. The IC₅₀ value obtained is that concentration of test compound required to inhibit, by 50%, the calcium-dependent release of [³H]dopamine. The lower the IC₅₀, the greater the agonistic activity; that is, the closer the value is to the

IC₅₀ obtained using melatonin, the more closely the compound is said to mimic melatonin activity.

Example 25 utilizes a similar assay with a modification to determine whether a test compound antagonizes the melatonin-induced inhibition of the release of [³H]dopamine from rabbit retina membranes. In this example, the test compound is added before the first period of electrical stimulation (S₁) and remains present until the end of the experiment. In this modification, melatonin at different concentrations is added before the second period of stimulation to ascertain whether the test compound, which is present during both S₁ and S₂ has the capacity to antagonize the action of melatonin. These assays are described in greater detail below.

EXAMPLE 23

15 Inhibition of 2-[¹²⁵I] iodomelatonin binding:

It has been shown that 2-[¹²⁵I]iodomelatonin binds with high affinity to a site exhibiting the pharmacological characteristics of a melatonin receptor in chicken retinal membranes. In this assay, i.e., in chicken retinal membranes, 2-[¹²⁵I]iodomelatonin selectively labels melatonin binding sites and shows no binding to alpha- and beta-adrenergic, serotonin, or dopamine sites, as agents that affect these sites do not compete for 2-[¹²⁵I]iodomelatonin binding. Dubocovich, et al., Proc. Nat'l. Acad. Sci. (USA), 84:3916 (1987). The test compounds were assayed for their possible melatonin agonist or antagonist-like activity, using an in vitro chicken retinal membrane preparation as described in Dubocovich, et al., Proc. Nat'l. Acad. Sci. (USA), 84:3916 (1987).

2-[¹²⁵I]iodomelatonin was diluted in Tris-HCl buffer (50 mM) with 0.01% bovine serum albumin. The compounds to be tested were dissolved in 1 mM HCl with 0.1% bovine serum albumin. Binding was initiated by addition of 220-μl aliquots of retinal membranes resuspended in the Tris-HCl buffer to tubes containing 20 μl of test compound or vehicle, and radioligand. The binding of 2-[¹²⁵I] iodomelatonin was routinely measured in duplicate after incubation at 0 °C for 5 hr in the dark. Reactions were terminated by addition of 5 ml of ice-cold Tris-HCl buffer, and the contents were immediately filtered through glass-fiber filters (Schleicher & Schuell no. 30) soaked in 0.5% (vol/vol) polyethylenimine solution. Each filter was washed twice with 5 ml of the cold buffer. Radioactivity was determined in a gamma counter. In a typical experiment, total 2-[¹²⁵I] iodomelatonin (67 pM) binding was 2542±361 cpm (n=3), and the non-specific binding defined with 3 μM 6-chloromelatonin was 350±35 cpm (n=3). The total radioactivity bound to filters represented 166±10 cpm (n=3).

The IC₅₀ is that concentration of a compound required to competitively inhibit the specific binding of [¹²⁵I]iodomelatonin by 50%. The results in Table 2 demonstrate that several compounds, in particular AH 001, AH 002, AH 013, and AH 017, AH 023, AH 024, AH 025, AH 026, AH 027, and AH 029 show high activity (low IC₅₀) in competitively inhibiting the binding of 2-[¹²⁵I]iodomelatonin to the melatonin binding site. This ability to competitively inhibit 2-[¹²⁵I]iodomelatonin demonstrates that these compounds interact with melatonin binding sites.

Table 2

Inhibition of 2-[¹²⁵ I]iodomelatonin Binding to Chicken Retinal Membranes	
Compound	IC ₅₀ (nM)
2-Iodomelatonin	2.5
6-Chloromelatonin	4.0
Melatonin	6.3
AH 001	860
AH 002	61
AH 005	20,000
AH 006	13,000
AH 007	>10,000
AH 008	>10,000
AH 010	>100,000
AH 011	20,000
AH 013	257
AH 014	4,620
AH 015	2,943
AH 016	>100,000
AH 017	682
AH 018	16,000
AH 019	>10,000
AH 020	2,908
AH 023	630
AH 024	250
AH 025	400
AH 026	250
AH 027	790
AH 028	25,000
AH 029	300

EXAMPLE 24Assay to Ascertain Which Test Compounds Mimic the Melatonin-Induced Inhibition of [³H]-dopamine Release from Rabbit Retina Membranes:

Melatonin and related indoles are capable of inhibiting the calcium-dependent release of [³H]dopamine from chicken retina and from rabbit retina. Dubocovich, J. Pharm. Exp. Ther. , 234:395-401 (1985). The amount of inhibition, by a test compound, of the calcium-dependent release of [³H]dopamine is therefore a useful measure of whether a compound is a melatonin agonist. In other words, a test compound is characterized as an agonist if its presence results in an inhibition of [³H]dopamine release.


Compounds inhibiting dopamine release may exert that affect by activating D-2 dopamine autoreceptors rather than melatonin receptors. Accordingly, all experiments were run in the presence of S-sulpiride (0.1 μM) to block D-2 dopamine autoreceptors present in the dopamine amacrine cells of retina.

IC₅₀ values are the concentration of test compound required to inhibit, by 50%, the calcium-dependent release of [³H]dopamine. Thus, the lower the IC₅₀, the greater the agonistic activity.

Assays were performed as described in Dubocovich, J. Pharm. Exp. Ther. , 234:395-401 (1985). Retinal tissues are incubated for 10 min at 37 °C in the presence of 0.1 μM [³H]dopamine. The tissue was then washed in Krebs' solution at 37 °C and transferred to individual cylindrical plastic tubes having a thin nylon

mesh on the bottom. These plastic tubes were then transferred to individual glass superfusion chambers containing platinum electrodes 30 mm apart. The tissue samples are superfused with Krebs' solution, starting at time equal zero until a leveling off of the spontaneous release of radioactivity occurred (about 60 mins). A time equal 60 mins S_1 is applied. Melatonin, or the test compound, is added 20 mins after S_1 and was present throughout the remainder of the experiment. The second stimulation (S_2) is applied 20 mins after the addition of melatonin or of the test compound.

Tritium release, i.e., release of [^3H]dopamine, was elicited by field stimulation at 3Hz, 20 mA, 2 msec duration. Field stimulations were applied in each experiment at either 60 (S_1) or 100 (S_2), min after the end of the incubation with [^3H]dopamine. Samples of the superfusate were collected before, during, and after the period of stimulation. At the end of each experiment, the retinal tissue contained in each chamber was solubilized and the tritium content remaining in the retinal tissue was determined by liquid scintillation counting.



In controls, when two periods of field stimulation are applied, at 60 (S_1) and 100 (S_2) min, the ratio of the amount of tritium released after S_2 to the amount released after S_1 , i.e., S_2/S_1 , is approximately one. That is, the same cumulative amount of radioactivity is released after each stimulation. However, if a drug, such as melatonin is added after the first stimulation, (but before the second stimulation), then the amount of radioactivity released after the second stimulation is significantly reduced and the S_2/S_1 ratio becomes less than one. Typically, the inhibition of the release of [^3H]dopamine, using 0.1 nM melatonin, is greater than half. If the S_2/S_1 is plotted as a function of the molar concentration of the amount of added melatonin, it is found that 40 pM melatonin will inhibit [^3H]dopamine release by 50%. Dubocovich, M.L., *J. Pharmacol. Exp. Ther.*, 246:902-904 (1988).

The inhibition of [^3H]dopamine overflow (the percentage of total tissue radioactivity released by field stimulation above the spontaneous release levels, expressed as the ratio S_2/S_1) is plotted as a function of the molar concentrations of the test compound (logarithmic scale). The IC_{50} values were determined graphically from these concentration effect curves (not shown) and are given in Table 3. These results demonstrate that compounds AH 001, AH 002, AH 011, AH 013, AH 015, AH 017, AH 025 and AH 027 are melatonin agonists. Test compounds that show no inhibition of the calcium-dependent release of ^3H -dopamine when added alone before S_2 , are assayed as melatonin receptor antagonists (See Example 25).

Table 3

Inhibition of [³ H]dopamine release from rabbit retina		
Compound	IC ₅₀ (nM) ^(a)	% Maximal Inhibition ^(b)
Melatonin	0.040	80
AH 001	1.4	80
AH 002	0.482	80
AH 005	52.2	50
AH 006	8.13	70
AH 007	10	40
AH 008	ND	ND
AH 009	ND	ND
AH 010	ND	ND
AH 011	1.3	80
AH 013	1.18	80
AH 014	7.94	80
AH 015	2.51	80
AH 016	>1,000	0
AH 017	0.063	75
AH 018	ND	ND
AH 019	ND	ND
AH 020	ND	ND
AH 023	NE	ND
AH 024	NE	ND
AH 025	4	70
AH 026	NE	ND
AH 027	13	65
AH 028	16	60
AH 029	10	70

a) IC₅₀ values represent the concentration of test compound necessary to inhibit the release of ³H-dopamine by 50%.

b) Percent maximal inhibition of ³H-dopamine release achieved with 1 μ M concentration of test compound.

ND = not determined

NE = no effect

EXAMPLE 25

Assay to Acertain Which Test Compounds Antagonize the Melatonin-Induced Inhibition of [³H]-dopamine Release from Rabbit Retina Membranes

Melatonin receptor antagonists are expected to prevent the inhibition of the calcium-dependent release of [³H]dopamine elicited by melatonin from rabbit retina. Dubocovich, M.L., J. Pharmacol. Exp. Ther. , 246:902 (1988).


Compounds that show competition for 2-[¹²⁵I]-iodomelatonin binding in chicken retinal membranes (Table 2) and show no inhibition of the calcium-dependent release of [³H]-dopamine when added alone before S₂ are tested for activity as melatonin receptor antagonists. (See Example 24).

To determine whether a compound is an antagonist of the melatonin-induced inhibition of [³H]dopamine release, the test compound is added before S₁ and remains present until the end of the experiment.

Melatonin (either 0.01, 0.1, 1.0, 10, or 100 nM) is then added after S_1 , but before S_2 .

Typically, the rabbit retina tissues pre-incubation with [^3H]-dopamine for 10 minutes is followed by a superfusion with Krebs solution at time equal zero. At time equal 20 minutes the test compound of interest is added. At time equal 60 minutes S_1 is applied. Melatonin is added 20 minutes after S_1 . S_2 applied 20 minutes after the addition of melatonin.

This relative time sequence is shown below:



For the control, no test compound is added and melatonin is added before S_2 ; this reduces the calcium-dependent release of [^3H]-dopamine resulting in an S_2/S_1 ratio of less than one. For example, in this particular experiment, using 0.1 nM melatonin gave an S_2/S_1 ratio of 0.52; while using 1 nM melatonin gave an S_2/S_1 ratio of 0.31.

As noted above, to determine whether a test compound has melatonin antagonist activity, the compound is added 40 minutes before S_1 and remains present throughout S_1 and S_2 . Approximately 20 minutes after S_1 , melatonin is added and remains present throughout S_2 . If a compound is an antagonist to melatonin activity, higher concentrations of melatonin are needed to attain the S_2/S_1 ratios obtained with melatonin in the absence of an antagonist. For test compounds AH 023, AH 024, and AH 026, S_2/S_1 ratios closer to unity were obtained (Table 4). These results demonstrate that these compounds do antagonize the melatonin-induced inhibition of [^3H]-dopamine release in rabbit retina.

The ratio S_2/S_1 can be plotted as a function of the molar concentrations of the test compound used (logarithmic scale). The concentration effect curves, obtained in the presence of 0.1 μM and 1 μM of either AH 023 or AH 024 (data not shown) show a shift to the right compared to the curve obtained with melatonin alone. In other words, the IC_{50} value obtained for melatonin alone is a lower molar concentration than obtained in the presence of the test compound, i.e., more melatonin is required to inhibit [^3H]-dopamine release in the presence of these antagonist compounds.

Table 4

Antagonism of melatonin-induced inhibition of [³ H]dopamine release from rabbit retina			
	S ₂ /S ₁		
Melatonin Concentration:	0.1 nM	1 nM	10 nM
Melatonin [†]	0.52±0.09 ^a (4)	0.31±0.06 ^a (6)	0.30±0.03(3)
AH 023 ^e			
0.01 μM	0.69±0.11(6)	0.30±0.04(6)	ND
0.1 μM	ND	0.73±0.04 ^c (6)	0.37±0.06(6)
1 μM	1.02±0.03 ^b (2)	0.92±0.07 ^c (2)	0.69±0.03 ^d (3)
AH 024 ^e			
0.1 μM	1.01±0.07 ^b (3)	0.76±0.01 ^c (6)	0.25±0.01(3)
1 μM	ND	0.91±0.06 ^c (3)	0.61±0.01 ^d (3)
AH 026 ^e			
0.1 μM	1.06±0.06 ^b (3)	0.91±0.05 ^c (3)	0.53±0.08 ^d (2)
1 μM	ND	1.08±0.03 ^c (3)	0.94±0.03 ^d (3)

a) $p < 0.05$ when compared with control (S_2/S_1 ratio = 0.98 ± 0.05 , $N = 14$).

b) $p < 0.05$ when compared with melatonin 0.1 nM.

c) $p < 0.05$ when compared with melatonin 1 nM.

d) $p < 0.05$ when compared with melatonin 10 nM.

e) test compounds in the concentrations indicated were added 40 minutes before S₁, and remained present through the experiment; melatonin (either 0.1 nM; 1 nM; or 10 nM; as indicated) was added 20 mins after S₁ $\phi^0 > \dagger$

π) μεωατοη η ψοηηθωδ δ τη ηο τεστ ψοηροξηηφ αφφεφ†

EXAMPLE 26

The compounds of the invention have melatonin agonist/antagonist activity and are expected to have utility in the treatment of disorders associated with circadian or seasonal rhythms and/or associated with abnormal melatonin levels in the body and/or associated with processes regulated by melatonin.

Melatonin has been implicated in the regulation of various neural and endocrine processes that are cued by the daily change in photoperiod. These include the regulation of seasonal effects on reproduction, body weight, metabolism, and coat color in photoperiodic mammals such as sheep and hamsters (Darrow, J.M., et al., J. Biol. Rhythms., 1:39-54 (1985); Tamarkin, L., et al., Science, 227:714-720 (1985); the control of circadian rhythms in birds and reptiles (Menaker, M., Vertebrate Circadian Systems, (Berlin:Springer-Verlag) 1-11 (1982); Underwood, H., et al., Physiol. Behav., 35:267-270 (1985)), and the modulation of retinal physiology (Iuvone, P.M., The retina: A Model for Cell Biology Studies, (London:Academic) 2:1-72 (1986); Besharse, J.C., Progress in Retinal Research, (Oxford:Pergamon) 1:81-124 (1982)). There is also evidence for photoperiodic and melatonin mediated control of cell cancer growth. Stanberry, et al., Endocrinol., 113:469-475 (1983). In addition, melatonin was shown to augment the immune response. Maestroni, et al., Clin. Exp. Immunol., 68:384-391 (1987).

Melatonin also has been implicated in a variety of brain functions and neuroendocrine changes, with a main site of action at the level of the hypothalamic-pituitary axis (Cardinali, Endocrinol. Rev., 2:327-346 (1981); and Minneman, et al., Life Sci., 17:1189-1200 (1975)). Abnormal plasma melatonin levels in humans has been associated with psychiatric illnesses (Jimerson, et al., Life Sci., 20:1501-1508 (1977); Smith, et al., J. Pharm Pharmacol., 31:246-248 (1979); and Ferrier, et al., Clin. Endocrinol., 17:181-187 (1982), with patients with pinealoma (Wetterberg, J. Neural Transm., 13:289-310 (1978)) and with breast

cancer (Tamarkin, et al., *Science*, 216:1003-1005 (1982)).

Melatonin may be effective in synchronizing disturbed circadian rhythms in mammals, including humans (Redman, J., et al., *Science*, 219:1089-1091 (1983); Cassone, V.M., et al., *J. Biol. Rhythms*, 1:219-229 (1986); Arendt, J., et al., *Br. Med. J.*, 292:1170 (1986)). For example, sleep/wake cycle disturbances, caused for example by jet lag (which results from rapid flight across several time zones) or caused by working on night shifts, and the subsequent desynchronization of circadian rhythms can be treated by administration of melatonin at scheduled times (Arendt, J., et al., *Br. Med. J.*, 292:1170 (1986)).

Changes in melatonin levels have also been shown at different stages of the menstrual cycle, with different seasons (Birau, N., *Melatonin-current status and perspectives*, (Oxford:Pergamon) 297-326 (1981); and Kauppila, A., et al., *J. Clin. Endocrinol. & Metab.*, 65:823-828 (1987)), and also with age (Iguchi, H., et al., *J. Clin. Endocrinol. & Metab.*, 55:27-29 (1982)). Changes in melatonin levels have been associated with various pathological states and the mode of action of various therapeutic agents (Birau, N., *Melatonin-current status and perspectives*, (Oxford:Pergamon) 297-326 (1981)). Certain neuroendocrine effects of melatonin in mammals, which include changes in gonadal function and pituitary hormone secretion, appear to be mediated to some extent by the action of melatonin on the brain.

Exposure of animals to light, which inhibits the synthesis of melatonin, or removal of the pineal gland are methods classically used to antagonize the physiological effects of endogenous melatonin (Darrow, et al., *J. Biol. Rhythms*, 1:39-54 (1985); and Tamarkin, et al., *Science*, 227:714-720 (1985)).

Recently, several groups of investigators have used bright light to treat chronobiologic disorders, e.g., seasonal affective disorders (SAD), sleep disorders, and symptoms such as drowsiness and fatigue associate with disturbances in sleep/wake cycles, in humans (Lewy, et al., *Proc. Soc. Exp. Biol. Med.*, 183:11-18 (1986); Lewy, et al., *Science*, 235:352-354 (1987); and Rosenthal, et al., *Arch. Gen. Psychiatric*, 41:72-80 (1984)). If the therapeutic effects of light are related to the suppression of melatonin secretion, melatonin receptor antagonists of the invention are expected to be useful in treating chronobiologic disorders involving changes in the pattern of melatonin secretion.

Administration of melatonin at scheduled times is effective in entraining and synchronizing circadian rhythms regulated primarily by the hypothalamic suprachiasmatic nuclei (Underwood, H., et al., *Physiol. Behav.*, 35:267-270 (1985); Redman, J., et al., *Science*, 219:1089-1091 (1983); and Cassone, V.M., et al., *J. Biol. Rhythms*, 1:219-229 (1986)). In addition, pineal melatonin appears to regulate several neuroendocrine functions such as sleep (Birkland, A.J., *Neuroendocrinology*, 34:126-131 (1982)), thyroid hormone secretion (Vriend, J., *Med. Hypotheses*, 4:376-387 (1978)) growth hormone release (Smythe, G.A., et al., *Nature (London)*, 244:230-231 (1973)) and intraocular pressure (Krauss, G., et al., *Inv. Ophthalm. Vis. Sci.*, Vol. 26, Abst. 49 (1988)) in humans. A melatonin receptor antagonist, by blocking the melatonin receptor in target tissues, would be expected to mimic the effects of light and to be more effective than pinealectomy in antagonizing the effects of endogenous melatonin inasmuch as this hormone is also secreted by extrapineal tissues (Cardinali, D.P., *Endocrinol. Rev.*, 2:327-346 (1981)).

It is also anticipated that compounds of the invention will be useful in the treatment of various psychiatric disorders related to altered melatonin function or influenced by melatonin and circadian rhythms, e.g., affective disorders (mania and depression), alcoholism, stress, epilepsy, convulsions, anxiety, tremors induced by idiopathic parkinsonism, and adventitious movements induced by L-dopa movements; and in the treatment of various neuroendocrine disorders related to altered melatonin function or influenced by melatonin and biological rhythms (e.g., peptic ulceration, psoriasis, hair growth, and body weight) particularly relating to regulation of reproductive maturation and function, e.g., idiopathic delayed puberty, sudden infant death (SID), premature labor, infertility, antifertility, premenstrual syndrome, and sexual dysfunction (i.e., impotence and decreased orgasmic activity). It is also expected that compounds of the invention can be used to manipulate the breeding cycles, body weight, coat color and oviposition in animals.

Further, it is anticipated that the compounds of the invention will find utility in localizing melatonin receptors. For example, compounds of the invention can be suitably labelled (e.g., with fluorescent groups or radioactive or non-radioactive isotopes) to serve as markers for both *in vitro* and *in vivo* analyses of for example, tissue distribution of melatonin receptors.

The foregoing illustrative examples relate to novel compounds having melatonin receptor agonist/antagonist activities, in particular substituted 2-amido tetralin derivatives; as well as to pharmaceutical preparations comprising such compounds; and methods for using these compounds as therapeutic and diagnostic reagents. While the present invention has been described in terms of specific compounds, compositions, and methods, it is understood that variations and modifications will occur to those skilled in the art upon consideration of the present invention.

For example, it is envisioned that various other substituted 2-amido-tetralin derivatives, will also be effective according to the present invention.


Although the preferred compounds are: 8-methoxy-2-acetamido-tetralin; 8-methoxy-2-propionamido-tetralin; 2-chloroacetamido-tetralin; 8-methoxy-2-n-butyramido-tetralin; 8-methoxy-2-cyclopropanecarbonylamido-tetralin; 8-methoxy-2-chloroacetamido-tetralin; 4-phenyl-2-acetamido-tetralin; 4-phenyl-2-propionamido-tetralin; 4-benzyl-2-acetamido-tetralin; and 4-phenyl-2-chloroacetamido-tetralin; and 4-benzyl-2-propionamido-tetralin; it is not intended to preclude others such as: a) 8-methoxy-4-phenyl-2-acetamido-tetralin; b) 8-methoxy-4-phenyl-2-propionamido-tetralin; c) 8-methoxy-4-benzyl-2-acetamido-tetralin; and d) 8-methoxy-4-benzyl-2-propionamido-tetralin, which, based on structure activity relationships are predicted to be very potent melatonin receptor antagonists (a, b) or potent melatonin receptor agonists (c, d); or any other effective substituted 2-amido-tetralin amide derivative from being included in the scope of the present invention.

Numerous modifications and variations in the invention as described in the above illustrative examples are expected to occur to those skilled in the art and consequently only such limitations as appear in the appended claims should be placed thereon.

Accordingly it is intended in the appended claims to cover all such equivalent variations which come within the scope of the invention as claimed.

Claims

1. Compounds of the formula:



wherein:

R₁ is hydrogen, halogen, amino, amido, a C₁-4 alkyl, alkoxy, or alkoxyaryl;

R₂ is hydrogen, hydroxyl, halogen, amino, amido, aryl, mono- or di- C₁-4 alkylamino, C₁-4 alkylaryl, or alkoxyaryl, or C₁-4 alkyl, alkenyl, alkyne, or alkoxy;

R₃ is hydrogen, aryl, C₁-4 alkylaryl, or C₁-4 alkyl, alkenyl, or alkyne;

R₄ is aryl, C₁-4 alkylaryl, or C₁-4 alkyl, haloalkyl, or cycloalkyl;

R₅ is hydrogen, hydroxyl, halogen, oxo, aryl, C₁-4 alkylaryl, or C₁-4 alkyl;

wherein aryl substituents of R₂, R₃, R₄, and R₅ may optionally be halogen, hydroxyl, amino, mono- or di- C₁-4 alkylamino, or C₁-4 alkyl or alkoxy substituted, provided that when R₁ is methoxy, and R₂, R₃, and R₅ are hydrogen, R₄ is not methyl.

2. The compounds according to claim 1 wherein said alkoxy is methoxy.

3. The compounds according to claim 1 provided that when said R₁ is alkoxy, R₂ is not alkoxy.

4. The compounds according to claim 1 wherein said R₄ is methyl, chloromethyl, ethyl, propyl, cyclopropyl, n-butyl, iso-butyl, phenyl, or benzyl.

5. The compounds according to claim 1 wherein said R₅ is phenyl or benzyl.

6. The compounds according to claim 1 which are:

8-methoxy-2-propionamido-tetralin;

2-chloroacetamido-tetralin;

8-methoxy-2-n-butyramido-tetralin;

8-methoxy-2-cyclopropanecarbonylamido-tetralin;

8-methoxy-2-chloroacetamido-tetralin;

4-phenyl-2-acetamido-tetralin;


4-phenyl-2-propionamido-tetralin;

4-benzyl-2-acetamido-tetralin;

4-phenyl-2-chloroacetamido-tetralin; and

4-benzyl-2-propionamido-tetralin.

7. Compounds of the formula:



wherein:

R₁ is hydrogen, halogen, amino, amido, a C₁₋₄ alkyl, alkoxy, or alkoxyaryl;

R₂ is hydrogen, hydroxyl, halogen, amino, amido, aryl, mono- or di- C₁₋₄ alkylamino, C₁₋₄ alkylaryl, or alkoxyaryl, or C₁₋₄ alkyl, alkenyl, alkynyl, alkoxy;

R₃ is hydrogen, aryl, C₁₋₄ alkylaryl or C₁₋₄ alkyl, alkenyl, or alkynyl;

R₄ is aryl, C₁₋₄ alkylaryl, or C₁₋₄ alkyl, haloalkyl, or cycloalkyl;


R₅ is hydrogen, hydroxyl, halogen, oxo, aryl, C₁₋₄ alkylaryl, or C₁₋₄ alkyl;

wherein aryl substituents of R₂, R₃, R₄, and R₅ may optionally be halogen, hydroxyl, amino, mono- or di- C₁₋₄ alkylamino, or C₁₋₄ alkyl or alkoxy substituted, provided that when R₁ is methoxy, and R₂, R₃ and R₅ are hydrogen, R₄ is not methyl.

8. A method as set forth in claim 7 wherein R₅ is hydrogen.

9. A method as set forth in claim 7 wherein R₅ is phenyl.

10. A method for treatment of disorders associated with abnormal melatonin activity in an organism, said method comprising the step of administering a therapeutically effective amount of a composition comprising a compound of the formula:



wherein:

R₁ is hydrogen, halogen, amino, amido, a C₁₋₄ alkyl, alkoxy, or alkoxyaryl;

R₂ is hydrogen, hydroxyl, halogen, amino, amido, aryl, mono- or di- C₁₋₄ alkylamino, C₁₋₄ alkylaryl, or alkoxyaryl, or C₁₋₄ alkyl, alkenyl, alkynyl, alkoxy;

R₃ is hydrogen, aryl, C₁₋₄ alkylaryl or C₁₋₄ alkyl, alkenyl, or alkynyl;

R₄ is aryl, C₁₋₄ alkylaryl, or C₁₋₄ alkyl, haloalkyl, or cycloalkyl;

R₅ is hydrogen, hydroxyl, halogen, oxo, aryl, C₁₋₄ alkylaryl, or C₁₋₄ alkyl;

wherein aryl substituents of R₂, R₃, R₄, and R₅ may optionally be halogen, hydroxyl, amino, mono- or di- C₁₋₄ alkyl or alkoxy substituted.

11. The method according to claim 10 wherein said compound is a compound selected from the group consisting of:

8-methoxy-2-propionamido-tetralin;

2-chloroacetamido-tetralin;

8-methoxy-2-n-butylamido-tetralin;

8-methoxy-2-cyclopropanecarbonylamido-tetralin;

8-methoxy-2-chloroacetamido-tetralin;

4-phenyl-2-acetamido-tetralin;


4-phenyl-2-propionamido-tetralin;

4-benzyl-2-acetamido-tetralin;

4-phenyl-2-chloroacetamido-tetralin; and

4-benzyl-2-propionamido-tetralin.

12. A pharmaceutical composition for use in treatment of disorders associated with abnormal melatonin activity in an organism, said composition comprising a pharmaceutically acceptable solvent, diluent, adjuvant, or carrier and as the active ingredient, from about 0.001 mg to about 100 mg per kg of body weight of said organism of a compound of the formula:




wherein:

- R₁ is hydrogen, halogen, amino, amido, a C₁₋₄ alkyl, alkoxy, or alkoxyaryl;
 R₂ is hydrogen, hydroxyl, halogen, amino, amido, aryl, mono- or di- C₁₋₄ alkylamino, C₁₋₄ alkylaryl, or alkoxyaryl, or C₁₋₄ alkyl, alkenyl, alkynyl, alkoxy;
 R₃ is hydrogen, aryl, C₁₋₄ alkylaryl or C₁₋₄ alkyl, alkenyl, or alkynyl;
 R₄ is aryl, C₁₋₄ alkylaryl, or C₁₋₄ alkyl, haloalkyl, or cycloalkyl;
 R₅ is hydrogen, hydroxyl, halogen, oxo, aryl, C₁₋₄ alkylaryl, or C₁₋₄ alkyl;
 wherein aryl substituents of R₂, R₃, R₄, and R₅ may optionally be halogen, hydroxyl, amino, mono- or di- C₁₋₄ alkylamino, or C₁₋₄ alkyl or alkoxy substituted.

13. A method for treatment of disorders associated with abnormal melatonin activity in an organism, said method comprising the step of administering a pharmaceutical composition according to claim 12.

14. In a method inhibiting melatonin function by administering a melatonin antagonist, the improvement comprising the step of administering an effective amount of a compound of the formula:




wherein:

- R₁ is hydrogen, halogen, amino, amido, a C₁₋₄ alkyl, alkoxy, or alkoxyaryl;
 R₂ is hydrogen, hydroxyl, halogen, amino, amido, aryl, mono- or di- C₁₋₄ alkylamino, C₁₋₄ alkylaryl, or alkoxyaryl, or C₁₋₄ alkyl, alkenyl, alkynyl, alkoxy;
 R₃ is hydrogen, aryl, C₁₋₄ alkylaryl or C₁₋₄ alkyl, alkenyl, or alkynyl;
 R₄ is aryl, C₁₋₄ alkylaryl, or C₁₋₄ alkyl, haloalkyl, or cycloalkyl;
 R₅ is hydrogen, hydroxyl, halogen, oxo, aryl, C₁₋₄ alkylaryl, or C₁₋₄ alkyl;
 wherein aryl substituents of R₂, R₃, R₄, and R₅ may optionally be halogen, hydroxyl, amino, mono- or di- C₁₋₄ alkylamino, or C₁₋₄ alkyl or alkoxy substituted.

15. The method according to claim 14 wherein said compound is selected from the group consisting of:

- 4-phenyl-2-acetamido-tetralin;
 4-phenyl-2-propionamido-tetralin, and
 4-phenyl-2-chloroacetamido-tetralin.

16. In a method for mimicking melatonin function by administering a melatonin agonist, the improvement comprising the step of administering an effective amount of a compound of the formula:



wherein:

R₁ is hydrogen, halogen, amino, amido, a C₁₋₄ alkyl, alkoxy, or alkoxyaryl;

R₂ is hydrogen, hydroxyl, halogen, amino, amido, aryl, mono- or di- C₁₋₄ alkylamino, C₁₋₄ alkylaryl, or alkoxyaryl, or C₁₋₄ alkyl, alkenyl, alkynyl, alkoxy;

R₃ is hydrogen, aryl, C₁₋₄ alkylaryl or C₁₋₄ alkyl, alkenyl, or alkynyl;

R₄ is aryl, C₁₋₄ alkylaryl, or C₁₋₄ alkyl, haloalkyl, or cycloalkyl;

R₅ is hydrogen, hydroxyl, halogen, oxo, aryl, C₁₋₄ alkylaryl, or C₁₋₄ alkyl;

wherein aryl substituents of R₂, R₃, R₄, and R₅ may optionally be halogen, hydroxyl, amino, mono- or di- C₁₋₄ alkylamino, or C₁₋₄ alkyl or alkoxy substituted.

17. The method according to claim 16 wherein said compound is selected from the group consisting of:

8-methoxy-2-propionamido-tetralin;

2-chloroacetamido-tetralin;

8-methoxy-2-n-butyramido-tetralin;


8-methoxy-2-cyclopropanecarbonylamido-tetralin;

8-methoxy-2-chloroacetamido-tetralin;

4-benzyl-2-acetamido-tetralin; and

4-benzyl-2-propionamido-tetralin.

18. A method for localizing melatonin receptors in a sample, said method comprising the steps of: incubating said sample with a compound of the formula:



wherein:

R₁ is hydrogen, halogen, amino, amido, a C₁₋₄ alkyl, alkoxy, or alkoxyaryl;

R₂ is hydrogen, hydroxyl, halogen, amino, amido, aryl, mono- or di- C₁₋₄ alkylamino, C₁₋₄ alkylaryl, or alkoxyaryl, or C₁₋₄ alkyl, alkenyl, alkynyl, alkoxy;

R₃ is hydrogen, aryl, C₁₋₄ alkylaryl or C₁₋₄ alkyl, alkenyl, or alkynyl;

R₄ is aryl, C₁₋₄ alkylaryl, or C₁₋₄ alkyl, haloalkyl, or cycloalkyl;

R₅ is hydrogen, hydroxyl, halogen, oxo, aryl, C₁₋₄ alkylaryl, or C₁₋₄ alkyl;

wherein aryl substituents of R₂, R₃, R₄, and R₅ may optionally be halogen, hydroxyl, amino, mono- or di- C₁₋₄ alkylamino, or C₁₋₄ alkyl or alkoxy substituted, and wherein said compound is labelled with a marker; detecting said marker; and

localizing said melatonin receptors in said sample.

19. A pharmaceutical composition for use in treatment of disorders associated with abnormal melatonin activity in an organism, said composition comprising as the active ingredient a substituted derivative of 2-amidotetralin having melatonin activity.

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 420 064 A3

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: **90118198.2**

(51) Int. Cl.⁵: **C07C 233/23, C07C 233/06,
C07C 233/74, C07C 233/60,
A61K 31/16**

(22) Date of filing: **21.09.90**

(30) Priority: **25.09.89 US 411675**

(43) Date of publication of application:
03.04.91 Bulletin 91/14

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

(88) Date of deferred publication of the search report:
26.02.92 Bulletin 92/09

(71) Applicant: **NORTHWESTERN UNIVERSITY
633 Clark Street
Evanston Illinois 60208(US)**

(72) Inventor: **Horn, Alan S.
Noordwijkgweg 3A
NL-9804 RA Noordhorn(NL)
Inventor: Dubocovich, Margarita L.
2608 Noyes
Evanston, Illinois 60201(US)**

(74) Representative: **Patentanwälte Grünecker,
Kinkeldey, Stockmair & Partner
Maximilianstrasse 58
W-8000 München 22(DE)**

(54) **Substituted 2-amidotetralins as melatonin agonists and antagonists.**

(57) The present invention relates generally to compounds having melatonin receptor activities and in particular to substituted 2-amidotetralin derivatives; to pharmaceutical preparations comprising such compounds; and to methods for using these compounds as therapeutic and diagnostic reagents.

EP 0 420 064 A3



European
Patent Office

EUROPEAN SEARCH REPORT

Application Number

EP 90 11 8198

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
X	CHEMICAL ABSTRACTS vol. 55, no. 23, 13 November 1961, column 23560, abstract no. 23560d, Columbus, Ohio, US; Z. HORII et al.: "Syntheses of oxytocics" & Yakugaku Zasshi 1961, vol. 81, pages 636-639 - - -	1,6-8	C 07 C 233/23 C 07 C 233/06 C 07 C 233/74 C 07 C 233/60 A 61 K 31/16
X	CHEMISCHE BERICHTE vol. 110, no. 2, 1977, pages 778-787, DE; A.-D. SCHWANGHART et al.: "Präparative Trennversuche an optisch aktiven Polyamiden" * page 780 * - - -	1,7,8	
X	CHEMICAL ABSTRACTS vol. 75, no. 3, 19 July 1971, page 398, column 1, abstract no. 19621y, Columbus, Ohio, US; V. GHISLANDI et al.: "Optical resolution and configuration of alpha-cyclohexylbenzylamine 1,2,3,4-tetrahydro-1-naphthylamine, and 1,2,3,4-tetrahydro-2-naphthylamine" & Farmaco, Ed. Sci. 1971, vol. 26, no. 5, pages 474-486 - - -	1,7,8	
P,X	EP-A-0 334 538 (THE UPJOHN COMPANY) * abstract; page 2; page 7, lines 40-55 * - - -	1-4,6-8	
X	EP-A-0 253 257 (BAYER AG) * page 14; claims 1,2,8 * - - -	1,4,7	TECHNICAL FIELDS SEARCHED (Int. Cl.5)
X	EP-A-0 286 516 (ADIR ET COMPAGNIE) * page 12, lines 1-28; abstract * - - -	1,4,7,8	C 07 C 233/00 A 61 K 31/00
X	EP-A-0 074 903 (SYNTHELABO) * page 13, lines 1-13; abstract * - - -	1,4,7,8	
X	FR-A-2 391 189 (DAIICHI SEIYAKU CO LTD.) * page 22, line 32; page 15, line 12 * - - -	1,2,7,8	
	-/-		
The present search report has been drawn up for all claims			
Place of search Berlin		Date of completion of search 11 November 91	Examiner RUFET J.M.A.
CATEGORY OF CITED DOCUMENTS X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons - - - - - &: member of the same patent family, corresponding document			



European
Patent Office

EUROPEAN SEARCH REPORT

Application Number

EP 90 11 8198

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
X	DE-A-2 022 746 (BOEHRINGER MANNHEIM GMBH) * pages 8,9 *	1	
X	GB-A-2 160 200 (SANDOZ LTD.) * claims 3,4,6 *	1	
X	AUSTRALIAN JOURNAL OF CHEMISTRY vol. 20, no. 3, March 1967, pages 601-604, AU; D.M. TEMPLE et al.: "Some new sympathomimetic amines" * page 602 *	1,4	
X	JOURNAL OF MOLECULAR STRUCTURE vol. 218, 1990, pages 81-86, Elsevier, Amsterdam-Oxford-New York-Tokyo; D.G. ANTONOVIC et al.: "Synthesis and spectroscopic study of some new N-naphtyl substituted 2-phenylacetamides" * page 83 *	1,7,8	
X	JOURNAL OF MEDICINAL CHEMISTRY vol. 32, no. 2, 1989, pages 478-486, American Chemical Society, US; Y. QIZ- HUANG et al.: "Conformationally Restricted and Confor- mationally Defined Tyramine Analogues as Inhibitors of Phenylethanolamine N-Methyltransferase" * abstract; pages 480,485, column right *	1,7,8	
X	JOURNAL OF MEDICINAL CHEMISTRY vol. 25, no. 2, 1982, pages 136-141, American Chemical Society, US; R.M. DE- MARINIS et al.: "Alpha-Adrenergic Agents. 2. Synthesis and alpha1-Agonist Activity of 2-Aminotetralins1,2" * page 137; abstract *	1,3	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
Place of search		Date of completion of search	Examiner
Berlin		11 November 91	RUFET J.M.A.
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention</p> <p>E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons &: member of the same patent family, corresponding document</p>			



European
Patent Office

EUROPEAN SEARCH REPORT

Page 3

Application Number

EP 90 11 8198

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
P,X	EP-A-0 378 456 (MERRELL DOW PHARMACEUTICALS INC.) * page 4 *	1,7,8	
A	WO-A-8 901 472 (NELSON RESEARCH & DEVELOPMENT CO.) * abstract; claims 1,5,12 *	1,10,12, 14	
			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
The present search report has been drawn up for all claims			
Place of search Berlin		Date of completion of search 11 November 91	Examiner RUFET J.M.A.
CATEGORY OF CITED DOCUMENTS X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons &: member of the same patent family, corresponding document			